

From the Spectrum to Inflation : An Inverse Formula for the General Slow-Roll Spectrum

Minu Joy^{1*} Ewan D. Stewart^{1,2†} Jinn-Ouk Gong¹ Hyun-Chul Lee¹

¹ *Department of Physics, KAIST, Daejeon, Republic of Korea*

² *Department of Physics and Astronomy, University of Canterbury,
Christchurch, New Zealand*

February 2, 2008

Abstract

We propose a general inverse formula for extracting inflationary parameters from the power spectrum of cosmological perturbations. Under the general slow-roll scheme, which helps to probe the properties of inflation in a model independent way, we invert the leading order, single field, power spectrum formula. We also give some physically interesting examples to demonstrate its wide applicability and illuminate its properties.

*minujoy@muon.kaist.ac.kr

†On sabbatical leave from Department of Physics, KAIST, Daejeon, Republic of Korea

1 Introduction

It is believed that the power spectra of both the cosmic microwave background radiation and the large scale structure of the observable universe have evolved from a common primordial spectrum of curvature perturbations produced during inflation [1]. Through the observation of these spectra we can study the properties of inflation. For example, results from the Wilkinson Microwave Anisotropy Probe [2] and Sloan Digital Sky Survey [3] provide a remarkable set of data, which places strong restrictions on models of inflation [4].

There have been many studies on the accurate evaluation of the power spectrum, provided that the inflationary parameters are given. But practically we observe the power spectrum first, and then seek an appropriate inflationary model from which the observed spectrum could result [5]. In the context of the standard slow-roll approximation, such inversion is straightforward [5]. However, the standard slow-roll approximation makes strong assumptions about the properties of inflation, which have not yet been confirmed observationally. Thus we are lacking a model independent way of extracting useful information about inflationary parameters from the power spectrum.

The general slow-roll approximation [6] was introduced to eliminate these unjustified assumptions of standard slow-roll, instead essentially only relying on the observed approximate scale invariance of the spectrum. See Refs. [7, 8] for some other alternatives to the standard slow-roll approximation. The one limitation of general slow-roll is that it does not cover cases where super-horizon effects are dominant [9]. Also, in this paper, we limit ourselves to the case of single field models of inflation, though the single field general slow-roll formulae can also be applied to some multi-field models of inflation [10]. These extensions will be treated thoroughly in a separate publication [11].

In this paper, we invert the leading order single field general slow-roll formula for the power spectrum [6] to obtain a formula, Eq. (7), for inflationary parameters in terms of the primordial power spectrum. See Ref. [12] for an alternative method of inversion that also does not rely on the standard slow-roll approximation. In Section 2 we give our inverse, in Section 3 we give some examples which illustrate the use and properties of our inverse, and in the Appendix we give some alternative forms for some of our formulae.

2 General slow-roll formulae

For single field inflationary models, it is convenient to express inflationary quantities in terms of

$$f = \frac{2\pi a \xi \dot{\phi}}{H}, \quad (1)$$

where $\xi = -\int \frac{dt}{a} = \frac{1}{aH} \left(1 - \frac{\dot{H}}{H^2} + \dots\right)$ is minus the conformal time [6, 13]. We think of f as a function of $\ln \xi$ so that $f' \equiv df/d \ln \xi$.

To leading order in the general slow-roll approximation, the spectrum can be expressed as [6]

$$\ln \mathcal{P}(k) = \int_0^\infty \frac{d\xi}{\xi} [-k\xi W'(k\xi)] \left[\ln \left(\frac{1}{f^2} \right) + \frac{2}{3} \frac{f'}{f} \right]. \quad (2)$$

There are a variety of other forms for this formula, some of which are given in the Appendix. The window function $-x W'(x)$ is given by [‡]

$$W(x) = \frac{3 \sin(2x)}{2x^3} - \frac{3 \cos(2x)}{x^2} - \frac{3 \sin(2x)}{2x} - 1. \quad (3)$$

It has the asymptotic behavior

$$\lim_{x \rightarrow 0} W(x) = \frac{2}{5}x^2 + \mathcal{O}(x^4), \quad (4)$$

the window property

$$\int_0^\infty \frac{dx}{x} [-x W'(x)] = 1, \quad (5)$$

and the degeneracy

$$\int_0^\infty \frac{dx}{x} [-x W'(x)] \frac{1}{x} = 0. \quad (6)$$

2.1 Inverse formula

Our inverse formula is

$$\ln \left(\frac{1}{f^2} \right) = \int_0^\infty \frac{dk}{k} m(k\xi) \ln \mathcal{P}, \quad (7)$$

where

$$m(x) = \frac{2}{\pi} \left[\frac{1}{x} - \frac{\cos(2x)}{x} - \sin(2x) \right]. \quad (8)$$

It has the asymptotic behavior

$$\lim_{x \rightarrow 0} m(x) = \frac{4}{3\pi}x^3 + \mathcal{O}(x^5) \quad (9)$$

and the window property

$$\int_0^\infty \frac{dx}{x} m(x) = 1. \quad (10)$$

It is straightforward to derive Eq. (7) from Eq. (2) using the key identity

$$\int_0^\infty \frac{dk}{k} m(k\zeta) W(k\xi) = \left(\frac{\zeta^3}{\xi^3} - 1 \right) \theta(\xi - \zeta), \quad (11)$$

where $\theta(x) = 0$ for $x < 0$ and $\theta(x) = 1$ for $x > 0$. An alternative form of the inverse formula is given in the Appendix.

[‡]Note that we define $W(x)$ with an extra -1 compared with our previous works [6].

3 Examples

3.1 Standard slow-roll approximation

In the context of standard slow-roll, the power spectrum has the form

$$\ln \mathcal{P} = \ln \mathcal{P}_\diamond + (n_\diamond - 1) \ln \left(\frac{k}{k_\diamond} \right) + \dots, \quad (12)$$

where $k_\diamond$ is some reference wavenumber. Applying our inverse formula, Eq. (7), gives

$$\ln \left(\frac{1}{f^2} \right) = \ln \mathcal{P}_\diamond + \alpha(n_\diamond - 1) - (n_\diamond - 1) \ln(k_\diamond \xi) + \dots, \quad (13)$$

where we have used Eq. (10) and

$$\int_0^\infty \frac{dx}{x} m(x) \ln x = \alpha, \quad (14)$$

where $\alpha = 2 - \ln 2 - \gamma \simeq 0.7296$. We see that Eq. (13) reproduces the standard slow-roll inverse, which is trivially obtained from the standard slow-roll formulae [13]

$$\ln \mathcal{P} = \ln \left(\frac{1}{f_\star^2} \right) - 2\alpha \frac{f'_\star}{f_\star} + 2 \frac{f'_\star}{f_\star} \ln(k \xi_\star) + \dots \quad (15)$$

and

$$n - 1 = 2 \frac{f'_\star}{f_\star} + \dots, \quad (16)$$

where $\xi_\star$ is an arbitrary evaluation point, usually taken to be around horizon crossing.

3.2 Power law

Consider a spectrum of the form

$$\ln \mathcal{P} = \ln \mathcal{P}_0 - A k^\nu \quad (17)$$

with $\nu > 0$. In the limit $\nu \ll 1$, this reproduces a significant subset of standard slow-roll spectra. However, we have in mind that instead it is A that is sufficiently small to give the required approximate scale invariance over observable scales. Besides being motivated by simplicity, such a form for the power spectrum is well motivated by inflationary model building [10].

Substituting into our inverse formula, Eq. (7), we obtain

$$\ln \left(\frac{1}{f^2} \right) = \ln \mathcal{P}_0 - A C(\nu) \xi^{-\nu}, \quad (18)$$

where

$$\begin{aligned} C(\nu) &= \int_0^\infty \frac{dx}{x} m(x) x^\nu \\ &= \frac{2^{1-\nu} \Gamma(2+\nu)}{\pi \nu(1-\nu)} \sin \left(\frac{\pi \nu}{2} \right) \quad \text{for } \nu < 1. \end{aligned} \quad (19)$$

For $\nu \geq 1$, the inverse diverges. This is to be expected as Eq. (6) shows that the forward formula, Eq. (2), is degenerate for $\nu = 1$.

For a better understanding of this degeneracy, we can start from an inflationary scenario [10] which leads to a power spectrum of the form of Eq. (17). In such a scenario, f is calculated to be

$$\ln \left(\frac{1}{f^2} \right) = \ln \left(\frac{1}{f_\infty^2} \right) - B \xi^{-\nu} \quad (20)$$

with $\nu > 0$. Then, calculating the power spectrum using Eq. (28), we obtain

$$\ln \mathcal{P} = \ln \left(\frac{1}{f_\infty^2} \right) - B D(\nu) k^\nu, \quad (21)$$

where

$$\begin{aligned} D(\nu) &= \int_0^\infty \frac{dx}{x} [-x w'(x)] x^{-\nu} \\ &= 2^\nu \frac{\Gamma(2-\nu)}{1+\nu} \cos \left(\frac{\pi\nu}{2} \right) \quad \text{for } \nu < 2. \end{aligned} \quad (22)$$

For $\nu < 1$ everything is straightforward and similar to the standard slow-roll case $\nu \ll 1$. For $\nu = 1$ we have $D(1) = 0$ so that the first order general slow-roll formula becomes degenerate. Thus we cannot expect an inverse to exist for $\nu \geq 1$. For $1 < \nu < 2$, the super-horizon tail of the window function $-x w'(x)$ dominates, but the general slow-roll formula, Eq. (28), still converges. For $\nu \geq 2$, the general slow-roll formula diverges indicating the need for super-horizon methods [9, 11].

Finally, from Eq. (18) or (21), we have

$$\ln \mathcal{P}_0 = \ln \left(\frac{1}{f_\infty^2} \right), \quad (23)$$

and noting that

$$C(\nu) D(\nu) = 1, \quad (24)$$

we see the consistency of Eqs. (17) and (18) with (20) and (21).

3.3 Linear potential with a sharp step

As our final example, we consider a linear potential of slope A with a sharp downward step at ϕ_s . See Ref. [14] for a more complete discussion of this example. The potential is

$$V(\phi) = V_0 [1 - A(\phi - \phi_s) - a \theta(\phi - \phi_s)], \quad (25)$$

where A determines the slope of the potential and a the height of the step. The potential is not smooth and we have a singularity at the position of the step. Therefore we cannot apply the standard slow-roll approximation since $\dot{\phi}$ is changing drastically at the step. But we can apply the general slow-roll approximation to calculate the power spectrum. We assume

$A \ll 1$ so that de Sitter space is a good approximation, and $aV_0 \ll A^2$ so that the step only generates a small departure from scale invariance. Then, we have

$$\ln \left(\frac{1}{f^2} \right) = \ln \left(\frac{V_0}{12\pi^2 A^2} \right) - 2 \frac{aV_0}{A^2} \left(\frac{\xi}{\xi_s} \right)^3 \theta(\xi_s - \xi). \quad (26)$$

Using Eq. (2), the power spectrum is

$$\ln \mathcal{P} = \ln \left(\frac{V_0}{12\pi^2 A^2} \right) + \frac{2}{3} \frac{aV_0}{A^2} k \xi_s W'(k \xi_s). \quad (27)$$

Now, using our inverse, Eq. (7), to invert this power spectrum we recover Eq. (26) by using Eq. (11). This again illustrates the power of the general slow-roll approach compared with the standard slow-roll approach.

Acknowledgements

We thank Jai-chan Hwang, Kenji Kadota, Misao Sasaki and Takahiro Tanaka for helpful discussions, and the Yukawa Institute for Theoretical Physics for hospitality while this work was in progress. JG is grateful to Jai-chan Hwang and Misao Sasaki for encouragement. This work was supported in part by ARCSEC funded by the Korea Science and Engineering Foundation and the Korean Ministry of Science, the Korea Research Foundation grant KRF PBRG 2002-070-C00022, Brain Korea 21, and an Erskine Fellowship of the University of Canterbury.

Appendix

In this appendix we give some alternative formulae for the spectrum and the inverse.

If $f_\infty \equiv \lim_{\xi \rightarrow \infty} f$ exists, as for example in the models of Ref. [10], then we have the following form

$$\ln \mathcal{P} = \ln \left(\frac{1}{f_\infty^2} \right) + \int_0^\infty \frac{d\xi}{\xi} [-k\xi w'(k\xi)] \ln \left(\frac{f_\infty^2}{f^2} \right), \quad (28)$$

where [§]

$$w(x) = \frac{\sin(2x)}{x} - \cos(2x) - 1. \quad (29)$$

It has the asymptotic behavior

$$\lim_{x \rightarrow 0} w(x) = \frac{2}{3}x^2 + \mathcal{O}(x^4), \quad (30)$$

the almost window property

$$\int_0^\infty \frac{dx}{x} [-x w'(x)] = 1 + \cos(2\infty), \quad (31)$$

[§]Note that we define $w(x)$ with an extra -1 compared with our previous works [6].

and the degeneracy

$$\int_0^\infty \frac{dx}{x} [-x w'(x)] \frac{1}{x} = 0. \quad (32)$$

It is related to $W(x)$ by

$$w(x) = W(x) + \frac{x}{3} W'(x), \quad (33)$$

and Eq. (11) gives

$$\int_0^\infty \frac{dk}{k} m(k\xi) w(k\xi) = -\theta(\xi - \zeta). \quad (34)$$

An alternative form for the spectrum with a particularly simple window function is

$$\ln \mathcal{P} = \int_0^\infty \frac{d\xi}{\xi} [-k\xi v'(k\xi)] \left[\ln \left(\frac{1}{f^2} \right) - 2 \frac{f'}{f} \right], \quad (35)$$

where

$$v(x) = \frac{\sin(2x)}{2x} - 1. \quad (36)$$

It has the asymptotic behavior

$$\lim_{x \rightarrow 0} v(x) = -\frac{2}{3}x^2 + \mathcal{O}(x^4), \quad (37)$$

window property

$$\int_0^\infty \frac{dx}{x} [-x v'(x)] = 1, \quad (38)$$

and is related to $w(x)$ by

$$w(x) = v(x) - x v'(x). \quad (39)$$

An alternative form for the inverse with a particularly simple window function is

$$\ln \left(\frac{1}{f^2} \right) = \int_0^\infty \frac{dk}{k} n(k\xi) \left[\ln \mathcal{P} + \frac{d \ln \mathcal{P}}{d \ln k} \right], \quad (40)$$

where

$$n(x) = \frac{1}{\pi} \left[\frac{1}{x} - \frac{\cos(2x)}{x} \right]. \quad (41)$$

It has the asymptotic behavior

$$\lim_{x \rightarrow 0} n(x) = -\frac{2}{\pi}x + \mathcal{O}(x^3), \quad (42)$$

window property

$$\int_0^\infty \frac{dx}{x} n(x) = 1, \quad (43)$$

and is related to $m(x)$ by

$$m(x) = n(x) - x n'(x). \quad (44)$$

References

- [1] A. A. Starobinsky, *Phys. Lett.* **B 117**, 175 (1982) ; A. Guth and S.-Y. Pi, *Phys. Rev. Lett.* **49**, 1100 (1982) ; J. M. Bardeen, P. J. Steinhardt and M. S. Turner, *Phys. Rev.* **D 28**, 679 (1983) ; H. Kodama and M. Sasaki, *Prog. Theor. Phys. Suppl.* **78**, 1 (1984)
- [2] C. L. Bennett et al., *Astrophys. J. Suppl.* **148**, 1 (2003) [astro-ph/0302207](#) ; D. N. Spergel et al., *Astrophys. J. Suppl.* **148**, 175 (2003) [astro-ph/0302209](#)
- [3] I. Zehavi et al., *Astrophys. J.* **571**, 172 (2002) [astro-ph/0106476](#) ; M. Tegmark et al., *Astrophys. J.* **606**, 702 (2004) [astro-ph/0310725](#)
- [4] M. Tegmark et al., *Phys. Rev.* **D 69**, 103501 (2004) [astro-ph/0310723](#)
- [5] E. J. Copeland, E. W. Kolb, A. R. Liddle and J. E. Lidsey, *Phys. Rev.* **D 48**, 2529 (1993) [hep-ph/9303288](#) ; E. J. Copeland, E. W. Kolb, A. R. Liddle and J. E. Lidsey, *Phys. Rev.* **D 49**, 1840 (1994) [astro-ph/9308044](#) ; J. E. Lidsey et al., *Rev. Mod. Phys.* **69**, 373 (1997) [astro-ph/9508078](#)
- [6] S. Dodelson and E. D. Stewart, *Phys. Rev.* **D 65**, 101301 (2002) [astro-ph/0109354](#) ; E. D. Stewart, *Phys. Rev.* **D 65**, 103508 (2002) [astro-ph/0110322](#) ; J. Choe, J.-O. Gong and E. D. Stewart, *JCAP* **07**, 012 (2004) [hep-ph/0405155](#) ; J.-O. Gong, *Class. Quant. Grav.* **21**, 5555 (2004) [gr-qc/0408039](#)
- [7] S. Habib, K. Heitmann, G. Jungman and C. Molina-Paris, *Phys. Rev. Lett.* **89**, 281301 (2002) [astro-ph/0208443](#) ; S. Habib, A. Heinen, K. Heitmann, G. Jungman and C. Molina-Paris, *Phys. Rev.* **D 70**, 083507 (2004) [astro-ph/0406134](#) ; S. Habib, A. Heinen, K. Heitmann and G. Jungman, *Phys. Rev.* **D 71**, 043518 (2005) [astro-ph/0501130](#)
- [8] J. Martin and D. J. Schwarz, *Phys. Rev.* **D 67**, 083512 (2003) [astro-ph/0210090](#)
- [9] S. M. Leach, M. Sasaki, D. Wands and A. R. Liddle, *Phys. Rev.* **D 64**, 023512 (2001) [astro-ph/0101406](#)
- [10] K. Kadota and E. D. Stewart, *JHEP* **07**, 013 (2003) [hep-ph/0304127](#) ; *JHEP* **12**, 008 (2003) [hep-ph/0311240](#)
- [11] H.-C. Lee, M. Sasaki, E. D. Stewart, T. Tanaka and S. Yokoyama, in preparation
- [12] S. Habib, K. Heitmann and G. Jungman, *Phys. Rev. Lett.* **94**, 061303 (2005) [astro-ph/0409599](#)
- [13] E. D. Stewart and J.-O. Gong, *Phys. Lett.* **B 510**, 1 (2001) [astro-ph/0101225](#)
- [14] J.-O. Gong, [astro-ph/0504383](#)